

TESTING THE VECTOR CONDENSATE MODEL OF ELECTROWEAK INTERACTIONS AT HIGH ENERGY HADRON COLLIDERS

G. Cynolter, E. Lendvai and G. Pócsik
Institute for Theoretical Physics,
Eötvös Lorand University, Budapest

Abstract

In the vector condensate model a doublet of vector fields plays the role of the Higgs doublet of standard model and the gauge symmetry is broken dynamically. This results in a theory surviving the test of radiative corrections provided the new charged and neutral vector particles B have masses of at least several hundred GeV's. In this note we show that while at the Tevatron the heavy B -particle production is too low, at LHC the yield is large and, for instance, the inclusive cross section of B^+B^- pairs is 51.5 (15.3) fb at $\sqrt{s} = 14$ TeV, $m_B = 400(500)$ GeV.

The vector condensate model employs the Lagrangian of the standard model of electroweak interactions but the usual scalar doublet is replaced by a doublet of vector fields B_μ [1] whose neutral component forms a condensate breaking the gauge symmetry dynamically and providing nonvanishing W, Z and vanishing photon masses. A quartic self-coupling of B_μ gives masses to the B -particles, as well as the interaction of B -pairs with fermions gives rise to fermion masses and embeds the Kobayashi-Maskawa mechanism [1].

Looking at the S parameter it follows that the B -particles are heavy, at $\Lambda = 1$ TeV the threshold is about $m_0 \simeq 400 - 500$ GeV for B^0 and $m_+ \simeq 200 - 350$ GeV for $B^\pm$ [2]. Higher Λ attracts higher minimum masses.

The allowed regions by S are tightened by the T parameter [3]. For example at $\Lambda = 1$ TeV, $m_0 = 400$ GeV the threshold for m_+ is increased to 630 GeV. In general, to each momentum scale there exists a range of B masses where the radiative corrections are suitably small by partial cancellations. This is why Λ cannot be too large as compared to B masses.

Recently it has been shown that pairs of heavy B-particles are copiously produced at high energy linear e^+e^- colliders [4]. In the present note we study the production of B-particles at the Tevatron and LHC. We show that producing heavy B-particles at LHC is very favourable having a large cross section while at Tevatron energy the production cross section cannot exceed (0.01-0.02) fb which is far below the discovery limit.

Since fermions are coupled very weakly to B-pairs [1] in the vector condensate model, producing B-pairs is expected to be more considerable from virtual γ and Z exchanges, that is we consider the Drell-Yan mechanism [5], $p(\bar{p}) \rightarrow B\bar{B} + X$ via quark-antiquark annihilation.

The Drell-Yan cross section for the above hadronic collisions can be written as [5,6]

$$\sigma(p(\bar{p}) \rightarrow B\bar{B} + X) = \int_{\tau_0}^1 d\tau \int_{\tau}^1 \frac{dx}{2x} \sum_i \sigma(q_i \bar{q}_i \rightarrow B\bar{B}) \cdot \left(f_i^1(x, \hat{s}) f_i^2(\tau/x, \hat{s}) + f_i^1(x, \hat{s}) f_i^2(\tau/x, \hat{s}) \right) \quad (1)$$

where x and τ/x are the parton momentum fractions, $\hat{s} = \tau s$ is the square of the centre of mass energy of $q_i \bar{q}_i$, s is the same for the hadronic initial state, $f_i^1(x, \hat{s})$ means the number distribution of i quarks in hadron 1 at the scale $\hat{s}$ and the sum runs over the quark flavours u,d,s,c. In the computation the MRS (G) fit program [7] was used for the parton distributions.

The angle integrated, colour averaged annihilation cross section $\sigma(q_i \bar{q}_i \rightarrow B\bar{B})$ is calculated to lowest order in the gauge couplings, and QCD corrections are neglected. We hope this approximation shows the order of magnitude of the cross section. For $B^0 \bar{B}^0$ final state the Z exchange is working, and $B^+ B^-$ pairs appear via $\gamma + Z$ exchange. This is because in the model the following Lagrangians relevant to the annihilation process emerge [1]:

$$\begin{aligned} L(B^0) &= \frac{ig}{2\cos\theta_W} \partial^\mu B^{(0)\nu+} \left(Z_\mu B_\nu^{(0)} - Z_\nu B_\mu^{(0)} \right) + h.c., \\ L(B^+ B^- Z) &= -\cos 2\theta_W \cdot L(B^{(0)} \rightarrow B^{(+)}) , \end{aligned} \quad (2)$$

where $B_\mu^{(0)}$ ($B_\mu^{(+)}$) denotes the field of neutral (charged) B-particles. At $q_j \bar{q}_j Z$ -vertex the usual coupling $ig\gamma_\mu(g_{Vj} + g_{Aj}\gamma_5)$ acts, here

$$\left. \begin{aligned} g_{Vj} &= \frac{1}{2} - \frac{4}{3}\sin^2\theta_W, \\ g_{Aj} &= \frac{1}{2} \end{aligned} \right\} j = u, c$$

$$\left. \begin{aligned} g_{Vj} &= -\frac{1}{2} + \frac{2}{3}\sin^2\theta_W, \\ g_{Aj} &= -\frac{1}{2} \end{aligned} \right\} j = d, s.$$
(3)

From (2) we get for $B^0 \bar{B}^0$ final states

$$\sigma(q_i \bar{q}_i \rightarrow B^0 \bar{B}^0) = \frac{1}{3} \frac{1}{256\pi} \left(\frac{g}{\cos\theta_W} \right)^4 (g_{Vi}^2 + g_{Ai}^2) \left(1 - \frac{4m_0^2}{\hat{s}} \right)^{3/2} \frac{\hat{s} + 8m_0^2}{4m_0^4}. \quad (4)$$

This is decreasing at high, increasing m_0 and for $\hat{s} \gg 4m_0^2$ it is proportional to $\hat{s}/m_0^4$ reflecting that the Lagrangian (2) is coming from a nonrenormalisable, effective model. The cross section of $B^+ B^-$ pairs can be expressed by (4) in the following way

$$\sigma(q_i \bar{q}_i \rightarrow B^+ B^-) = \sigma(q_i \bar{q}_i \rightarrow B^0 \bar{B}^0; m_0 \rightarrow m_+) \cdot \frac{1}{g_{Vi}^2 + g_{Ai}^2} \cdot \left[(g_{Vi}^2 + g_{Ai}^2) \cos^2 2\theta_W + 2Q_{q_i} g_{Vi} \sin^2 2\theta_W \cos 2\theta_W + 4Q_{q_i}^2 \sin^2 2\theta_W \right]. \quad (5)$$

The individual terms are due to Z exchange, $\gamma - Z$ interference and γ exchange.

We have calculated various distributions of $B\bar{B}$ pairs for $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV and for pp collisions at $\sqrt{s} = 14$ TeV, assuming $m_B = 400, 500, 600$ GeV. Typically, at the Tevatron no notable result can be presented, however, increasing the energy to LHC, we get sizable cross sections. Also the yield of $B^+ B^-$ is larger than that of $B^0 \bar{B}^0$. As an example we show in Fig.1 the τ -distribution of $B^+ B^-$ pairs at LHC, $m_+ = 400$ GeV. $\frac{d\sigma}{d\tau}$ is sharply peaked after threshold ($4m_+^2/s$) and insignificant from about $\tau = 0.1$, that is for higher invariant masses of $B^+ B^-$. Fig.2 shows $\frac{\partial^2 \sigma}{\partial p_T \partial y}|_{y=0}$ as the function of the transverse momentum p_T of B^+ at vanishing rapidity y of $B^+ B^-$ for LHC, $m_+ = 400$ GeV.

For the total cross section (1) we obtain

$$\begin{aligned}
\sigma_{\text{TeV}} &= 0.020(0.0145)fb \text{ for } B^+B^-(B^0\overline{B}^0), m_B = 400\text{GeV}, \\
\sigma_{LHC} &= 51.46; 15.30; 5.65fb \text{ (44.35; 13.54; 4.90fb)} \\
&\text{for } B^+B^-(B^0\overline{B}^0), m_B = 0.4; 0.5; 0.6\text{TeV}.
\end{aligned} \tag{6}$$

For instance, at an expected integrated luminosity of 10^5pb^{-1} one gets about 5100 B^+B^- pairs of $m_+ = 0.4$ TeV at LHC per annum.

In conclusion, we have shown that heavy B-particle pairs have a large inclusive cross section due to $q\overline{q}$ annihilation at LHC in a suitable mass range making the detection of $B^{+,0}$ at LHC possible.

This work is supported in part by OTKA I/7, No. 16248.

References

- [1] G. Pócsik, E. Lendvai and G. Cynolter, Acta Phys. Pol. **B24**, 1495 (1993); G.Cynolter, E. Lendvai and G. Pócsik, in Electroweak Symmetry Breaking, World Scientific, 1995 (Ed. F. Csikor and G. Pócsik) p. 85.
- [2] G. Cynolter, E. Lendvai and G. Pócsik, Mod. Phys. Lett. **A10**, 2193 (1995).
- [3] G. Cynolter, E. Lendvai and G. Pócsik, Mod. Phys. Lett. **A9** , 1701 (1994)
- [4] G. Cynolter, E. Lendvai and G. Pócsik, Acta Phys. Pol. **B26**, 921 (1995).
- [5] S.D. Drell and T.M. Yan, Phys. Rev. Lett. **25**, 316 (1970); Ann. Phys. (N.Y) **66**, 578 (1971).
- [6] E. Eichten, I. Hinchliffe, K. Lane and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984).
- [7] A.D. Martin, W.J. Stirling and R.G. Roberts, Phys. Lett **B 354**, 155 (1995).

Figure Captions

Fig. 1: The differential τ -distribution of B^+B^- pairs at LHC, $m_+ = 400$ GeV.

Fig. 2: $\frac{\partial^2 \sigma}{\partial p_T \partial y} \Big|_{y=0}$ at LHC, $m_+ = 400$ GeV. p_T denotes the transverse momentum of B^+ and y is the rapidity of B^+B^- .

Fig.1.




Fig.2.

